

# Metodi Matematici della Meccanica Quantistica

## Assignment 3

To be handed in on **Wednesday, November 13, 2023, before 20:59** via email  
(scanned or L<sup>A</sup>T<sub>E</sub>X) to [ngoc.nguyen@unimi.it](mailto:ngoc.nguyen@unimi.it).

### Problem 1: Weyl criterion (10 points)

Let  $X$  be a Banach space,  $A : D \subset X \rightarrow X$  an operator and  $\lambda \in \mathbb{C}$ .

Prove that: If there exists a Weyl sequence, i. e., a sequence  $(x_n)_{n \in \mathbb{N}}$  in  $D$  with  $\|x_n\| = 1$  for all  $n \in \mathbb{N}$  and  $\|(A - \lambda)x_n\| \rightarrow 0$  as  $n \rightarrow \infty$ , then  $\lambda \in \sigma(A)$ .

### Problem 2: Coulomb potential (5+5 points)

a. Suppose that  $V \in L^2 + L^\infty(\mathbb{R}^3)$ .

Show that the  $L^2$ -part can be made arbitrarily small, i. e., for every  $\varepsilon > 0$  there exist  $V_2^\varepsilon \in L^2(\mathbb{R}^3)$  and  $V_\infty^\varepsilon \in L^\infty(\mathbb{R}^3)$  such that  $V = V_2^\varepsilon + V_\infty^\varepsilon$  and  $\|V_2^\varepsilon\|_2 < \varepsilon$ .

b. Show that the Coulomb potential, defined by  $V(x) := \frac{1}{|x|}$  for  $x \neq 0$  and  $V(0) := 0$ , is in  $L^2 + L^\infty(\mathbb{R}^3)$ .

### Problem 3: Sobolev inequalities (5+5 points)

a. Assume that for functions defined on  $\mathbb{R}^n$  a Sobolev inequality of the following form holds: There exists a constant  $C_{n,p,q}$  such that for all  $f$  we have

$$\|f\|_{L^q} \leq C_{n,p,q} \|\nabla f\|_{L^p} .$$

Given  $n$  and  $p$ , consider a rescaling  $f_\lambda(x) = f(\lambda x)$  by a parameter  $\lambda > 0$  to determine the only possible exponent  $q$  for which this can hold.

b. Let  $u \in H^m(\mathbb{R}^n)$  with  $m > n/2$  (the Sobolev space such that  $u$  has  $m$  weak derivatives in  $L^2$ ). Use the Fourier transform to show that  $u \in L^\infty(\mathbb{R}^n)$  and

$$\|u\|_{L^\infty} \leq C_{m,n} \|u\|_{H^m} , \tag{1}$$

where the constant  $C_{m,n}$  does not depend on  $u$ .

**Problem 4: On the Existence Proof for SCUGs (6+6 points)**

Let  $A : D \subset \mathcal{H} \rightarrow \mathcal{H}$  be a self-adjoint operator in a Hilbert space  $\mathcal{H}$ .

In the lecture we defined  $B_m := im(A + im)^{-1}$  ( $m \in \mathbb{Z}$ ),  $A_m := B_m A B_{-m}$ , and

$$U_m(t) := e^{-iA_m t} := \sum_{k \in \mathbb{N}} \frac{1}{k!} (-itA_m)^k .$$

- a. Show that  $U_m(t)$  is a SCUG for every  $m \in \mathbb{N}$ . Then show that the limit of  $U_m(t)\varphi$  ( $m \rightarrow \infty$ ) exists for all  $\varphi \in D$  and all  $t \in \mathbb{R}$ . Why does  $U(t) := s\text{-}\lim_{m \rightarrow \infty} U_m(t)$  exist?
- b. Show that  $U(t)$  is a SCUG. Then show that its generator is  $A$ .

**Problem 5: Resolvent of the Laplacian (8 points)**

Let  $H_0 = -\Delta$  as an operator with domain  $H^2(\mathbb{R}^3)$  in  $L^2(\mathbb{R}^3)$ . Show that for  $\varphi \in L^2(\mathbb{R}^3)$  and  $\kappa > 0$  we have

$$\left( (H_0 + \kappa^2)^{-1} \varphi \right) (x) = \frac{1}{4\pi} \int_{\mathbb{R}^3} \frac{e^{-\kappa|x-y|}}{|x-y|} \varphi(y) dy .$$

*Hint:* Recall that the Fourier transform turns a multiplication operator into convolution with the inversely transformed function  $\check{f}$ :

$$(\mathcal{F}^{-1} T_f \mathcal{F} \varphi) (x) = (2\pi)^{-n/2} \int_{\mathbb{R}^n} \check{f}(x-y) \varphi(y) dy .$$

Then use spherical coordinates and the residue theorem of complex analysis (if necessary, look it up!) to calculate  $\check{f}$ .